

JMS (ISSN 1682-4474) is an International, peer-reviewed scientific journal that publishes original article in experimental & clinical medicine and related disciplines such as molecular biology, biochemistry, genetics, biophysics, bio-and medical technology. JMS is issued eight times per year on paper and in electronic format.

For further information about this article or if you need reprints, please contact:

Dr. Mansoor Keshavarz
Department of Physiology,
School of Medicine,
Medical Sciences/University of
Tehran, P.O. Box 14155-6447,
Tehran, Iran

Tel/Fax: +9821-66419484

J. Med. Sci., 8 (2): 162-169
15th February, 2008

The Influence of Temperature Alterations on Eccentric Contraction-Induced Isometric Force and Desmin Loss in Rat Medial Gastrocnemius Muscle

¹Behnoosh Vasaghi-Gharamaleki, ²Mansoor Keshavarz,
³Shahryar Gharibzadeh, ⁴Masoud Sotodeh, ⁵Hamidreza Marvi,
⁶Javad Mosayebnejad and ¹Ismail Ebrahimi Takamjani

In this study isolated perfused rat muscle was used to examine the direct effect of temperature changes on the eccentric contraction-induced force and desmin loss. The left medial gastrocnemius muscle was separated and the entire lower limb was transferred into a prewarmed (35°C) organ bath. Temperature was adjusted to 31 or 39°C before and during eccentric contractions. Maximal isometric force and desmin loss were measured after 15 isometric or eccentric contractions. According to our data, organ bath temperature changes before or during eccentric contractions had no significant effect on force loss. However, a strong correlation between desmin loss and temperature changes before ($r = 0.93$, $p < 0.05$) and during ($r = 0.87$, $p < 0.05$) eccentric contractions was observed. Present results suggest that cooling before or during eccentric contractions may decrease desmin loss.

Key words: Eccentric contraction, isolated perfused muscle, isometric force loss, desmin

¹Department of Sport Medicine, School of Rehabilitation Sciences,
Iran University of Medical Sciences, Tehran, Iran

²Department of Physiology, School of Medicine,
Medical Sciences/University of Tehran, Iran

³Neuromuscular System Laboratory, Faculty of Biomedical Engineering,
Amirkabir University of Technology, Tehran, Iran

⁴Department of Pathology, Shariaty Hospital,
Medical Science/University of Tehran, Tehran, Iran

⁵School of Mechanical Engineering, Sharif University of Technology, Tehran, Iran

⁶Department of Mechanical Engineering,
Iran University of Science and Technology, Tehran, Iran

INTRODUCTION

In many situations, contracted muscle may elongate. It is a kind of contraction called eccentric or lengthening contraction. These contractions occur during activities such as downhill running, walking downstairs or lowering a weight. Unaccustomed eccentric muscle contractions result in muscle damage (Allen, 2001). Magnitude of the injury was closely related to the magnitude of the muscle strain on the fibers (Lieber and Fridén, 2002).

The typical feature of muscle damage are high plasma Creatine Kinase (CK) activity, strength loss, range of motion loss, swelling and Delay Onset Muscle Soreness (DOMS) (Evans *et al.*, 2002). The timing of these changes suggests that the muscle weakness might be a primary consequence of the muscle damage (Armstrong *et al.*, 1991). The histological feature of muscle damage is probably the extensive sarcomere disruption and Z-disk streaming. Desmin is the main intermediate filament protein in skeletal and heart muscles. In mature skeletal muscle, desmin filaments encircle and interlink myofibrils at the level of the Z disks and connect them to the plasma membrane thus aligning the myofibrils (Price, 1984). It is shown that eccentric contractions produce an early activation of the ubiquitous calpains 1 and 2 whose *in vitro* substrates include desmin and titin (Belcastro, 1993).

Losses of force and desmin have been observed very soon (within minutes) (Lieber and Fridén, 2002; Fridén and Lieber, 2001). For this reason and because studies on isolated muscle can analyze force and desmin loss but not pain or tenderness, the focus of this article is on the early muscle weakness and desmin staining following eccentric muscle contractions.

The effect of temperature changes on the eccentric exercise-induced muscle damage has been investigated in different studies (Evans *et al.*, 2002; Warren *et al.*, 2002; Nosaka *et al.*, 2004; Brock Symons *et al.*, 2004), but effect of temperature alternations on ultrastructural changes after eccentric contractions have not studied to date. It has been shown that heating may be effective in reducing muscle injury symptoms such as swelling (Evans *et al.*, 2002; Safran *et al.*, 1988). Increased temperature can be achieved actively through exercise, or passively, through the use of heat modalities such as ultrasound, short-wave diathermy and warm-water immersion. Cooling also has direct and indirect effects on muscle performance (Frank, 2001). There are limited reports about the effect of cooling on eccentric contraction-induced muscle damage (Paddon-Jones and Quigley, 1997; Nosaka *et al.*, 2004).

The enzymatic activity is affected by temperature. Meanwhile the eccentric contraction heat production is low (Allen, 2001; Constable *et al.*, 1997). On the other hand, the ultrastructural evidence of muscle damage has been related to a loss of desmin (Fridén and Lieber, 1998,

2001). Therefore, in our study, the force and desmin loss were chosen to investigate whether they are affected by the changes in temperature and/or the time of application of the temperature changes.

MATERIALS AND METHODS

Experiments were performed in accordance with the criteria outlined in the Guide for the Care and Use of Laboratory Animals (Copyright 1996 by the National Academy of Sciences; <http://books.nap.edu/readingroom/books/labrats/>). The protocol of the present experiment was approved by the institutional ethics committee of Medical Sciences/University of Tehran. Male Sprague Dawley rats ($n = 47$) weighting between 270 and 300 g (19 to 21 weeks) were used. All animals were maintained at a constant temperature ($22 \pm 0.5^\circ\text{C}$) with 12/12 light/dark cycles and with free access to food and water.

Surgery: The animals were anesthetized with sodium pentobarbital (50 mg kg^{-1} , intra-peritoneally) and placed supine on the operating table. The skin was totally removed from the entire left lower limb. Femoral artery was exposed and cannulated with heparinized (200 U/cc) PE-10 catheter. To restrict perfusion to the lower portion of the limb, the catheter was advanced as far as possible toward the popliteal space. The muscle was perfused at 6 mL min^{-1} with modified Krebs Henseleit solution (containing in mM: NaCl; 118.5, KCl; 4.7, CaCl_2 ; 2.5, MgSO_4 ; 1.2, KH_2PO_4 ; 1.2, NaHCO_3 ; 25, glucose; 11) by a peristaltic pump (Meredos GmbH, Germany). The perfusate was gassed with 95% O_2 and 5% CO_2 (final pH 7.4) and its temperature was maintained at 35°C and continuously recorded by a digital thermometer at the point before entering the catheter into the artery.

Medial gastrocnemius muscle has its own separate nerve and vessels and can be isolated without injury to its structure (Rijkelijhuizen *et al.*, 2005). It can also be perfused via femoral artery cannula. After cannulation, the thigh muscles and femur bone were cut. The bone was clamped in horizontal position. Medial gastrocnemius muscle was exposed by removing the semitendinosus and biceps femoris muscles as previously described by Rijkelijhuizen *et al.* (2005). Then the femur was clamped in a vertical position, whereas the medial gastrocnemius was fixed horizontally. Left lower limb and medial gastrocnemius muscle were transferred into water-jacket perfusion chamber. At the end of surgery, animals were killed with high doses of sodium pentobarbital.

Temperature selection: Since the perfusion fluid temperature was kept constant during the study and the eccentric contraction heat production was low (Allen,

2001; Constable *et al.*, 1997), the organ bath temperature may be the main factor determining muscle temperature. The temperatures selected for use were based on the following reasons. First, superficial and deep muscle temperatures of rats at rest are normally 35.5-37.2°C (Delp *et al.*, 1999). Similar values have been observed for human, but temperatures in the 30-35°C range are not uncommon (Ranatunga *et al.*, 1987; Kenny *et al.*, 2003). Second, different studies have shown 1 to 4°C temperature rise during active and passive warm-up (Draper *et al.*, 1998; Ashton *et al.*, 1998; Wirth *et al.*, 1998; Kenny *et al.*, 2003; Draper *et al.*, 1999) and icing decrease muscle temperature at least 5°C (Nosaka *et al.*, 2004). Third, although muscle temperature during exercise exceeds 40°C, experiments were not performed at temperature above 39°C because of concerns about muscle viability in our set-up. Therefore, organ bath temperature was adjusted to 35±4°C.

Experimental protocol. An important issue in all studies of eccentric muscle damage is to distinguish between the reduction in force caused by fatigue and that caused by the eccentric contractions. For this reason, it is important to compare the eccentric exercise against an isometric (or concentric) control and to design the study so that the reduction in force after the isometric series is minimal (Allen, 2001). Thus this study was conducted in isometric and eccentric sections. In each section the experimental protocol was the same.

Groups: After transportation of each muscle into chamber, tissue was allowed adapting for 45 min (Organ bath temperature was adjusted to 35°C) and then the experimental protocol continued in 5 different ways:

- **G35 (control):** The temperature was set at 35°C during the entire experiment.
- **G31B:** The temperature was set at 31°C for 30 min before eccentric contractions and then kept at 35°C during eccentric contractions.
- **G31D:** The temperature was set at 31°C during eccentric contractions for 30 min.
- **G39B:** The temperature was set at 39°C for 30 min before eccentric contractions and then kept at 35°C during eccentric contractions.
- **G39D:** The temperature was set at 39°C during eccentric contractions.

Muscle perfusion was continued for 30 min at 35°C after contraction period in all groups.

Muscle perfusion: To ascertain complete perfusion, we have conducted a pilot study in which medial gastrocnemius muscle samples were frozen after about

160 min perfusion (with or without isometric contractions) and non-perfusion period. Then transverse slices of 2 mm thickness were made and incubated in 1% triphenyltetrazoliumchloride (TTC) in phosphate buffer solution (pH 7.4) for 30 min at 37°C. To enhance the contrast between stained and unstained tissues, slices were immersed in 10% formalin. Stained slices were viewed under a light stereoscope. Brick red stained tissues were taken as viable, whereas pale or white ones were considered as necrotic.

Electrical stimulation: Medial gastrocnemius muscle has its own nerve and vessel branches and can be easily stimulated (Rijkkelijkhuizen *et al.*, 2005). To isolate the effect of temperature on nerve conduction, the muscle was stimulated directly. Stimulating electrodes (0.2 mm diameter platinum wire) were implanted into each end of the muscle (out of tissue sampling area) and connected to an electrical stimulator (Harvard 6002, Harvard Apparatus, Holliston, Massachusetts USA). Supramaximal stimulation (15-20 V) consisted of 100-Hz stimulus (square pulse, 0.5 msec width) with a train duration of 0.7 sec.

Eccentric contractions: The Achilles tendon together with a piece of calcaneal bone was connected to a force transducer (MLT50 ADInstruments, Australia) mounted on a servomotor. Acceleration, velocity, start length, onset of movement, stimulation algorithm, stimulation frequency and duration of the muscle contractions were controlled by a computer. Force and length outputs from the servomotor were sampled at 1 kHz.

The optimum length (L_0) of the medial gastrocnemius was defined as the muscle length at which active force was maximal. Eccentric contractions were performed for a range of 8 mm below to 4 mm over L_0 . For each eccentric contraction, the muscle was first placed at $L_0 - 8$ and isometrically activated until tension stabilized (300 msec) and then lengthening change was imposed at a rate of 17 mm sec⁻¹ for 0.7 sec⁻¹. Thus total duration of each contraction was approximately 1 sec. After each contraction, the muscle was allowed to recover at $L_0 - 8$ mm for 2 min to minimize the effect of fatigue. Fifteen eccentric contractions were carried out. Recording was performed by Power Lab System (4SP, ADInstruments, Australia). Force data were also saved on a compact disk for later analysis.

Isometric force loss: Isometric force measuring has been the most widely used method of determining muscle function after eccentric exercise (Warren *et al.*, 1999). To determine force loss due to eccentric contractions, maximal isometric force was measured before and 10 min after isometric or eccentric contractions. Supramaximal isometric stimulation (15-20 V) was applied with a

frequency of 100 Hz and train duration of 300 msec. Percentage change in maximal isometric force from pre- to post-eccentric contractions was used to indicate the magnitude of the eccentric contraction-induced force loss. In all groups, maximal isometric force was recorded at 35°C.

Desmin staining: Muscle sections were immunostained with antidesmin (monoclonal mouse anti-human clone D33, from Dako, Glostrup, Denmark) to evaluate the structural integrity of the cytoskeletal network (Fridén and Lieber, 1998). All muscle tissue samples were obtained from predetermined location of medial gastrocnemius muscle. Since, perfused muscles with isometric contractions gave typical cross-sectional patterns at different temperatures and did not show desmin loss only eccentric contractions groups were studied in this stage. For semiquantitative analysis (Koh and Escobedo, 2004), three independent observers blindly examined cross sections from muscles exposed to eccentric contractions at different temperatures. For each section, each observer counted the number of fibers that demonstrated overtly altered staining patterns after eccentric contractions (reduced staining of desmin protein, Fig. 1), normalized this number to the total number of fibers in the section




Fig. 1: Eccentric contraction-induced desmin loss. A, desmin staining of muscle sample after isometric contractions, that doesn't show any desmin loss. B, muscle sample after eccentric contractions in which desmin staining is lost. Bar 100 μ m

and normalized values were averaged between the three observers. Desmin staining was calculated from the following formula:

$$\text{Desmin staining} = 100 - \text{Desmin loss}$$

Statistical analysis: Data were expressed as mean \pm SEM. Comparison between isometric force before and after eccentric and isometric contractions was done by Student's paired t test. Groups were compared by one-way ANOVA and Tukey's post hoc test. Pearson's correlation coefficient (r) was used for desmin loss in different temperatures. All statistical analyses were performed using SPSS software (SPSS, Chicago, IL, USA). p-values less than 0.05 were considered statistically significant in all groups.

RESULTS

TTC staining: TTC staining was used to ensure muscle perfusion. It didn't reveal any necrotic area in perfused muscles (with and without isometric contractions) after 160 min, but in non-perfused samples the necrotic areas and ischemic slices were distinguishable.

Isometric force loss: As mentioned before, percentage change in maximal isometric force from pre - to post-contractions was used to indicate the magnitude of the contraction-induced force loss (Warren *et al.*, 1999). By using Student's paired t-test, isometric contractions showed no significant force loss ($p > 0.05$) but, exposure to eccentric contractions was associated with a significant reduction in maximum isometric force ($p < 0.05$). However, organ bath temperature changes before or during isometric or eccentric contractions had no statistically significant effect on force loss (Fig. 2, $p > 0.05$).

Desmin loss: The observations were blindly done (inter reliability, ICC = 0.801). Isometric contractions had no effect on desmin pattern but in eccentric groups, there was a significant difference ($p < 0.05$) in desmin loss among different groups (Fig. 3).

As it shown in Fig. 3, all eccentric groups had significant difference comparing with control group (G35). The most desmin loss was seen in G39B. Temperature increment before eccentric contractions induced more desmin loss than during it. However, there is no significant difference in desmin loss when organ bath temperature decreased to 31°C before or during eccentric contractions.

Pearsons correlation was performed to see if any relationship existed between desmin loss and temperature changes (Fig. 4). According to our data, there were a strong correlation between desmin loss and temperature




Fig. 2: Effect of organ bath temperature changes on contraction-induced isometric force loss. There were no significant differences among isometric (n = 3-4) or eccentric (n = 6-7) groups, $p > 0.05$. G35, the temperature was set at 35°C during the entire experiment. G31B, before contractions, the temperature was set at 31°C for 30 min. G31D, during contractions, the temperature was set at 31°C for 30 min. G39B, before contractions, the temperature was set at 39°C for 30 min. G39D, during contractions, the temperature was set at 39°C for 30 min. Values with the same letter are not significantly different. Values are mean±SEM




Fig. 3: Effect of organ bath temperature on desmin loss. G35, the temperature was set at 35°C during the entire experiment. G31B, before contractions, the temperature was set at 31°C for 30 min. G31D, during contractions, the temperature was set at 31°C for 30 min. G39B, before contractions, the temperature was set at 39°C for 30 min. G39D, during contractions, the temperature was set at 39°C for 30 min. Values with the same letter are not significantly different. Values are mean±SEM. $p < 0.05$

changes before ($r = 0.93$, $p < 0.05$) and during ($r = 0.89$, $p < 0.05$) eccentric contractions (Fig. 4A, B).

DISCUSSION

We have investigated the effect of temperature changes on force and desmin loss resulting from eccentric contractions of isolated perfused medial gastrocnemius muscle. In this study, there was a strong effect of temperature on the desmin loss caused by a bout of eccentric contractions. In contrast, there was no effect of




Fig. 4: Scatter plots of desmin loss vs organ bath temperature. There is a strong correlation between desmin loss and organ bath temperature before (A) and during (B) eccentric contractions

temperature changes on eccentric contraction-induced force loss.

Since *in vivo* changes in muscle temperature can increase sympathetic nerve activity via metaboreflexes and mechanoreflexes which result in systemic effects (Ray and Gracey, 1997; Ray *et al.*, 1997), it is difficult to identify the direct effect of temperature. On the other hand, in isolated muscle, the eccentric contraction-induced force loss can be attributed to hypoxia (Warren *et al.*, 2002). Moreover, although muscle's architecture is an important determinant of its force generating capacity (Blazevich and Sharp, 2005), the orientation of muscle fibers in force transmission may be disturbed due to muscle dissection in isolated muscle preparation. In the present study to eliminate above factors we have used isolated perfused medial gastrocnemius muscle to evaluate the direct effect of temperature changes on the eccentric contraction-induced muscle damage.

Since working muscle is affected by its perfusion, to ensure muscle full perfusion and oxygen supply, TTC staining, a fast and cheap method for detecting tissue injury by measuring dehydrogenase activity, was used (Yaoita *et al.*, 1993; Goldlust *et al.*, 1996). TTC staining in present study showed complete medial gastrocnemius muscle perfusion.

The eccentric contractions-induced force loss after unaccustomed eccentric contractions is a long-lasting phenomenon and complete recovery can take place in more than 1 month (Howell *et al.*, 1993). The exact cause of the eccentric contraction-induced force loss is unknown (Allen, 2001; Friden and Lieber, 2001; Evans *et al.*, 2002; Proske and Allen, 2005). According to our results, exposure to eccentric contractions was associated with a significant reduction in maximum isometric force but, there were no significant differences in eccentric contraction-induced isometric force loss and the time of temperature change among the isolated perfused skeletal muscles tested at three temperature points (Fig. 2). The same results have been obtained in human studies when temperature changes were applied before eccentric contractions (Evans *et al.*, 2002; Tiidus *et al.*, 2002; Nosaka *et al.*, 2004; Brock Symons *et al.*, 2004). For example, Nosaka *et al.* (2004) found that passive warm-up (short wave diathermy) and cool-down (ice pack) before eccentric contractions had no effect on eccentric contraction-induced isometric force loss. Since, our data showed that temperature changes had no effect on isometric force loss in isolated perfused skeletal muscle as reported in human studies, we concluded that the muscle temperature control systems may have no important role on eccentric contraction-induced isometric force loss.

Warren *et al.* (2002) reported the effect of temperature changes on isometric force loss in isolated muscle. They changed the temperature during eccentric contractions and found strong positive relationship between temperature changes and eccentric contraction-induced isometric force loss in isolated muscle. The contrast between the two studies may arise from more simulation of normal condition in present study such as the preservation of tendon-bone complex or full perfusion of muscle in our set-up

As mentioned earlier, ultrastructural evidence of muscle damage has been related to a loss of desmin (Fridén and Lieber, 1998, 2001), the major intermediate filament protein in striated muscle (Price, 1984). Desmin loss after eccentric contraction is extremely rapid and occurs within 5 min. This dramatic and rapid desmin loss, which does not occur after either isometric or concentric contraction, points to some type of enzymatic hydrolysis or protein phosphorylation as a likely mechanism rather than gene regulation, which requires much more time (Lieber and Fridén, 2002). It was shown that exercise-induced muscle damage produces an early activation of the ubiquitous calpains 1 and 2 whose *in vitro* substrates include desmin and titin (Belcastro, 1993). Meanwhile, ultrastructural evidence of muscle damage has been related to a loss of desmin (Fridén and Lieber, 1998, 2001).

According to present results, as the organ bath temperature was increased in eccentric contractions groups, desmin loss was increased too. On the other hand, isometric contractions at different temperatures had no effect on desmin staining. In eccentric groups, if the temperature changes were applied before eccentric contractions, a stronger correlation was seen than when temperature changes were applied during eccentric contractions (Fig. 4). To our knowledge, there are no published data on this filed and according to these results we suggest that the mechanism(s) that underlying the desmin loss act better at higher temperatures especially when the organ bath temperature changes were applied before the eccentric contractions. Unfortunately, we did not examine the calpain enzyme activity in this study and the tissue samples of this study are not suitable for enzymatic analysis. Although the specific mechanisms involved in this response remain to be determined, our results suggest that the mechanism(s) that underlying the desmin loss act better at higher temperatures especially when the organ bath temperature changes were applied before the eccentric contractions. Accordingly, it is suggested to examine the effect of temperature changes on calpains enzyme activity in eccentric contractions in the separate study in the future.

During the past 20 years the structural changes that occur in response to eccentric exercise have been interpreted as muscle injury but Yu *et al.* (2003) findings have suggested an active remodeling process in response to eccentric exercise in humans. It seems that after a focal initial loss of some myofibrillar proteins, an addition of new sarcomeres into pre-existing sarcomeres can be observed. Although Yu *et al.* (2003) did not see desmin loss 1 h, 2-3 and 7-8 days after eccentric exercise in human samples, in several other studies on both human and animals, desmin loss were seen (Fridén and Lieber, 1998; Beaton *et al.*, 2002; Peters *et al.*, 2003; Crameri *et al.*, 2007). In our study desmin loss was occurred and it was temperature sensitive.

Training of muscle by using eccentric contractions is popular among athletes and bodybuilders, as they have found this to be an effective strategy to increase muscular strength. Understanding of the effect of muscle temperature or temperature modalities on eccentric contractions will allow us to use the most appropriate temperature.

In conclusion, the present study may provide some insight into the factors that affect the desmin loss in eccentric contractions. Present findings suggest that desmin loss in spite of the force loss is temperature sensitive and additional research on the use of thermal modalities is encouraged to provide better rationale for application of them.

ACKNOWLEDGMENTS

This study was supported by a grant from the Research Council of Medical Sciences/University of Tehran, Tehran, Iran. The authors gratefully acknowledge Dr. Shadan F., Dr. Djahanguiry B., Dr. Mirzaii I and Mr. Sohanaki H. for reviewing the manuscript.

REFERENCES

- Allen, D.G., 2001. Eccentric muscle damage: Mechanisms of early reduction of force. *Acta Physiol. Scand.*, 171 (3): 311-319.
- Armstrong, R.B., G.L. Warren and J.A. Warren, 1991. Mechanisms of exercise-induced muscle fiber injury. *Sports Med.*, 12 (3): 184-207.
- Ashton, D.F., D.O. Draper and J.W. Myrer, 1998. Temperature rise in human muscle during ultrasound treatments using flex-all as a coupling agent. *J. Athl. Train.*, 33 (2): 136-140.
- Beaton, L.J., M.A. Tarnopolsky and S.M. Phillips, 2002. Contraction-induced muscle damage in humans following calcium channel blocker administration. *J. Physiol.*, 544 (Pt 3): 849-859.
- Belcastro, A.N., 1993. Skeletal muscle calcium-activated neutral protease (calpain) with exercise. *J. Applied Physiol.*, 74 (30): 1381-1386.
- Blazevich, A.J. and N.C.C. Sharp, 2005. Understanding muscle architectural adaptation: Macro-and micro-level research. *Cells Tissue Organs*, 181 (1): 1-10.
- Brock Symons, T., J.L. Clasey, D.R. Gater and J.W. Yates, 2004. Effects of deep heat as a preventative mechanism on delayed onset muscle soreness. *J. Strength. Cond. Res.*, 18 (1): 155-161.
- Constable, J.K., C.J. Barclay and C.L. Gibbs, 1997. Energetics of lengthening in mouse and toad skeletal muscles. *J. Physiol.*, 505 (Pt 1): 205-215.
- Cramer, R.M., P. Aagaard, K. Qvortrup, H. Langberg, J. Olesen and M. Kjaer, 2007. Myofiber damage in human skeletal muscle: Effects of electrical stimulation versus voluntary contraction. *J. Physiol.*, 583 (1): 365-380.
- Delp, M.D., C. Duan, C.A. Ray and R.B. Armstrong, 1999. Rat hindlimb muscle blood flow during level and downhill locomotion. *J. Applied Physiol.*, 86 (2): 564-568.
- Draper, D., St. Harris, S. Schulthies, E. Durrant, K.L. Knight and M. Ricard, 1998. Hot-Pack and 1-MHZ ultrasound treatments have an additive effect on muscle temperature increase. *J. Athl. Train.*, 33 (1): 21-24.
- Draper, D.O., K. Knight, T. Fujiwara and J.C. Castel, 1999. Temperature change in human muscle during and after pulsed short-wave diathermy. *J. Orthop. Sports Phy. Ther.*, 29 (1): 13-18; discussion 19-22.
- Evans, R.K., K.L. Knight, D.O. Draper and A.C. Parcell, 2002. Effects of warm-up before eccentric exercise on indirect markers of muscle damage. *Med. Sci. Sports Exerc.*, 34 (12): 1892-1899.
- Frank, S.M., 2001. Consequence of hypothermia; Focus on: Perioperative hypothermia. *Curr. Anesth. Crit. Care*, 12 (2): 79-86.
- Fridén, J. and R.L. Lieber, 1998. Segmental muscle fiber lesions after repetitive eccentric contractions. *Cell Tissue Res.*, 293 (1): 165-171.
- Fridén, J. and R.L. Lieber, 2001. Eccentric exercise-induced injuries to contractile and cytoskeletal muscle fiber components. *Acta Physiol. Scand.*, 171 (3): 321-326.
- Goldlust, E.J., R.P. Paczynski, Y.Y. He, C.Y. Hsu and M.P. Goldberg, 1996. Automated measurement of infarct size with scanned images of triphenyltetrazolium chloride-stained rat brains. *Stroke*, 27 (9): 1657-1662.
- Howell, J.N., G. Chleboun and R. Conatser, 1993. Muscle stiffness, strength loss, swelling and soreness following exercise-induced injury in humans. *J. Physiol.*, 464 (1): 183-196.
- Kenny, G.P., F.D. Reardon, W. Zaleski, M.L. Reardon, F. Haman and M.B. Ducharme, 2003. Muscle temperature transients before, during and after exercise measured using an intramuscular multisensor probe. *J. Applied Physiol.*, 94 (6): 2350-2357.
- Koh, T.J. and J. Escobedo, 2004. Cytoskeletal disruption and small heat shock protein translocation immediately after lengthening contractions. *Am. J. Physiol. Cell Physiol.*, 286 (3): C713-C722.
- Lieber, R.L. and J. Fridén, 2002. Mechanisms of muscle injury gleaned from animal models. *Am. J. Phys. Med. Rehabil.*, 81 (Suppl. 11): S70-S79.
- Nosaka, K., K. Sakamoto, M. Mewton and P. Sacco, 2004. Influence of pre-exercise muscle temperature on responses to eccentric exercise. *J. Athl. Train.*, 39 (2): 132-137.
- Paddon-Jones, D. and B.M. Quigley, 1997. Effect of cryotherapy on muscle soreness and strength following eccentric exercise. *Int. J. Sports Med.*, 18 (8): 588-593.
- Peters, D., I.A. Barash, M. Burdi, P.S. Yuan, L. Mathew, J. Fridén and R. Lieber, 2003. Asynchronous functional, cellular and transcriptional changes after a bout of eccentric exercise in rat. *J. Physiol.*, 553 (Pt 3): 947-957.

- Price, M.G., 1984. Molecular analysis of intermediate filament cytoskeleton-putative load-bearing structure. *Am. J. Physiol.*, 246 (4 Pt 2): H566-H572.
- Proske, U. and T.J. Allen, 2005. Damage to skeletal muscle from eccentric exercise. *Exerc. Sport Sci. Rev.*, 33 (2): 98-104.
- Ranatunga, K.W., B. Sharpe and B. Turnbull, 1987. Contractions of a human skeletal muscle at different temperatures. *J. Physiol.*, 390 (1): 383-395.
- Ray, C.A. and K.H. Gracey, 1997. Augmentation of exercise-induced muscle sympathetic nerve activity during muscle heating. *J. Applied Physiol.*, 82 (6): 1719-1725.
- Ray, C.A., K.M. Hume, K. H. Gracey and E.T. Mahoney, 1997. Muscle cooling delays activation of the muscle metaboreflex in humans. *Am. J. Physiol.*, 273 (5 Pt 2): H2436-2441.
- Rijkelijkhuizen, J.M., G.C. Baan, A. de Haan, C.J. de Ruiter and P.A. Huijting, 2005. Extramuscular myofascial force transmission for *in situ* rat medial gastrocnemius and plantaris muscles in progressive stages of dissection. *J. Exp. Biol.*, 208 (Pt 1): 129-140.
- Safran, M.R., W.E.J. Garrett, A.V. Seaber, R.R. Glisson and B.M. Ribbeck, 1988. The role of warm up in muscular injury prevention. *Am. J. Sports Med.*, 16 (2): 123-129.
- Tiidus, P.M., J. Cort, S.J. Woodruff and P. Bryden, 2002. Ultrasound treatment and recovery from eccentric-exercise-induced muscle damage. *J. Sport Rehabil.*, 11 (4): 305-314.
- Warren, G.L., D.A. Lowe and R.B. Armstrong, 1999. Measurement tools used in the study of eccentric contraction-induced injury. *Sports Med.*, 27 (1): 43-59.
- Warren, G.L., C.P. Ingalls and R.B. Armstrong, 2002. Temperature dependency of force loss and Ca (2+) homeostasis in mouse EDL muscle after eccentric contractions. *Am. J. Physiol. Regul. Integr. Comp. Physiol.*, 282 (4): R1122-R1132.
- Wirth, V.J., B.L. Van Luen, D. Mistry, E. Saliba and F.C. McCue, 1998. Temperature changes in deep muscles of humans during upper and lower extremity exercise. *J. Athl. Train.*, 33 (3): 211-215.
- Yacita, H., A.J. Fischman, H.W. Strauss, T. Saito, E. Sato and Y. Maruyama, 1993. Uridine: A marker of myocardial viability after coronary occlusion and reperfusion. *Int. J. Cardiac Imaging*, 9 (4): 273-280.
- Yu, J.G., D.O. Fürst and L.E. Thomell, 2003. The mode of myofibril remodelling in human skeletal muscle affected by DOMS induced by eccentric contractions. *Histochem. Cell. Biol.*, 119 (5): 383-393.